



ECOSYSTEMS AND HUMAN WELL-BEING

Synthesis



MILLENNIUM ECOSYSTEM ASSESSMENT

Suggested citation:

Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-being: Synthesis*.
Island Press, Washington, DC.

Copyright © 2005 World Resources Institute

All rights reserved under International and Pan-American Copyright Conventions. No part of this book
may be reproduced in any form or by any means without permission in writing from the copyright holder:
World Resources Institute, 10 G Street NE, Suite 800, Washington, DC 20002.

ISLAND PRESS is a trademark of The Center for Resource Economics.

Library of Congress Cataloging-in-Publication data.

Ecosystems and human well-being : synthesis / Millennium Ecosystem Assessment.

p. cm. – (The Millennium Ecosystem Assessment series)

ISBN 1-59726-040-1 (pbk. : alk. paper)


1. Human ecology. 2. Ecosystem management. I. Millennium Ecosystem Assessment (Program) II. Series.

GF50.E26 2005

304.2–dc22

2005010265

British Cataloguing-in-Publication data available.

Printed on recycled, acid-free paper 

Book design by Dever Designs

Manufactured in the United States of America

CONTENTS

Foreword	ii
Preface	v
Reader's Guide	x
Summary for Decision-makers	1
Finding 1: Ecosystem Change in Last 50 Years	2
Finding 2: Gains and Losses from Ecosystem Change	5
Finding 3: Ecosystem Prospects for Next 50 Years	14
Finding 4: Reversing Ecosystem Degradation	18
Key Questions in the Millennium Ecosystem Assessment	25
1. How have ecosystems changed?	26
2. How have ecosystem services and their uses changed?	39
3. How have ecosystem changes affected human well-being and poverty alleviation?	49
4. What are the most critical factors causing ecosystem changes?	64
5. How might ecosystems and their services change in the future under various plausible scenarios?	71
6. What can be learned about the consequences of ecosystem change for human well-being at sub-global scales?	84
7. What is known about time scales, inertia, and the risk of nonlinear changes in ecosystems?	88
8. What options exist to manage ecosystems sustainably?	92
9. What are the most important uncertainties hindering decision-making concerning ecosystems?	101
Appendix A. Ecosystem Service Reports	103
Appendix B. Effectiveness of Assessed Responses	123
Appendix C. Authors, Coordinators, and Review Editors	132
Appendix D. Abbreviations, Acronyms, and Figure Sources	136
Appendix E. Assessment Report Tables of Contents	137

PREFACE

The Millennium Ecosystem Assessment was carried out between 2001 and 2005 to assess the consequences of ecosystem change for human well-being and to establish the scientific basis for actions needed to enhance the conservation and sustainable use of ecosystems and their contributions to human well-being. The MA responds to government requests for information received through four international conventions—the Convention on Biological Diversity, the United Nations Convention to Combat Desertification, the Ramsar Convention on Wetlands, and the Convention on Migratory Species—and is designed to also meet needs of other stakeholders, including the business community, the health sector, nongovernmental organizations, and indigenous peoples. The sub-global assessments also aimed to meet the needs of users in the regions where they were undertaken.

The assessment focuses on the linkages between ecosystems and human well-being and, in particular, on “ecosystem services.” An ecosystem is a dynamic complex of plant, animal, and microorganism communities and the nonliving environment interacting as a functional unit. The MA deals with the full range of ecosystems—from those relatively undisturbed, such as natural forests, to landscapes with mixed patterns of human use, to ecosystems intensively managed and modified by humans, such as agricultural land and urban areas. Ecosystem services are the benefits people obtain from ecosystems. These include *provisioning services* such as food, water, timber, and fiber; *regulating services* that affect climate, floods, disease, wastes, and water quality; *cultural services* that provide recreational, aesthetic, and spiritual benefits; and *supporting services* such as soil formation, photosynthesis, and nutrient cycling. (See Figure A.) The human species, while buffered against environmental changes by culture and technology, is fundamentally dependent on the flow of ecosystem services.


The MA examines how changes in ecosystem services influence human well-being. Human well-being is assumed to have multiple constituents, including the *basic material for a good life*, such as secure and adequate livelihoods, enough food at all times, shelter, clothing, and access to goods; *health*, including feeling well and having a healthy physical environment, such as clean air and access to clean water; *good social relations*, including social cohesion, mutual respect, and the ability to help others and provide for children; *security*, including secure access to natural and other resources, personal safety, and security from natural and human-made disasters; and *freedom of choice and action*, including the opportunity to achieve what an individual values doing and being. Freedom of choice and action is influenced by other constituents of well-being (as well as by other factors, notably education) and is also a precondition for achieving other components of well-being, particularly with respect to equity and fairness.

The conceptual framework for the MA posits that people are integral parts of ecosystems and that a dynamic interaction exists between them and other parts of ecosystems, with the changing human condition driving, both directly and indirectly, changes in ecosystems and thereby causing changes in human well-being. (See Figure B.) At the same time, social, economic, and cultural factors unrelated to ecosystems alter the human condition, and many natural forces influence ecosystems. Although the MA emphasizes the linkages between ecosystems and human well-being, it recognizes that the actions people take that influence ecosystems result not just from concern about human well-being but also from considerations of the intrinsic value of species and ecosystems. Intrinsic value is the value of something in and for itself, irrespective of its utility for someone else.

The Millennium Ecosystem Assessment synthesizes information from the scientific literature and relevant peer-reviewed datasets and models. It incorporates knowledge held by the private sector, practitioners, local communities, and indigenous peoples. The MA did not aim to generate new primary knowledge, but instead sought to add value to existing information by collating, evaluating, summarizing, interpreting, and communicating it in a useful form. Assessments like this one apply the judgment of experts to existing knowledge to provide scientifically credible answers to policy-relevant questions. The focus on policy-relevant questions and the explicit use of expert judgment distinguish this type of assessment from a scientific review.

Figure A. LINKAGES BETWEEN ECOSYSTEM SERVICES AND HUMAN WELL-BEING

This Figure depicts the strength of linkages between categories of ecosystem services and components of human well-being that are commonly encountered, and includes indications of the extent to which it is possible for socioeconomic factors to mediate the linkage. (For example, if it is possible to purchase a substitute for a degraded ecosystem service, then there is a high potential for mediation.) The strength of the linkages and the potential for mediation differ in different ecosystems and regions. In addition to the influence of ecosystem services on human well-being depicted here, other factors—including other environmental factors as well as economic, social, technological, and cultural factors—influence human well-being, and ecosystems are in turn affected by changes in human well-being. (See Figure B.)




Source: Millennium Ecosystem Assessment

ARROW'S COLOR	ARROW'S WIDTH
Potential for mediation by socioeconomic factors	Intensity of linkages between ecosystem services and human well-being
Low	Weak
Medium	Medium
High	Strong

Figure B. MILLENNIUM ECOSYSTEM ASSESSMENT CONCEPTUAL FRAMEWORK OF INTERACTIONS BETWEEN BIODIVERSITY, ECOSYSTEM SERVICES, HUMAN WELL-BEING, AND DRIVERS OF CHANGE

Changes in drivers that indirectly affect biodiversity, such as population, technology, and lifestyle (upper right corner of Figure), can lead to changes in drivers directly affecting biodiversity, such as the catch of fish or the application of fertilizers (lower right corner). These result in changes to ecosystems and the services they provide (lower left corner), thereby affecting human well-being. These interactions can take place at more than one scale and can cross scales. For example, an international demand for timber may lead to a regional loss of forest cover, which increases flood magnitude along a local stretch of a river. Similarly, the interactions can take place across different time scales. Different strategies and interventions can be applied at many points in this framework to enhance human well-being and conserve ecosystems.



Five overarching questions, along with more detailed lists of user needs developed through discussions with stakeholders or provided by governments through international conventions, guided the issues that were assessed:

- What are the current condition and trends of ecosystems, ecosystem services, and human well-being?
- What are plausible future changes in ecosystems and their ecosystem services and the consequent changes in human well-being?
- What can be done to enhance well-being and conserve ecosystems? What are the strengths and weaknesses of response options that can be considered to realize or avoid specific futures?
- What are the key uncertainties that hinder effective decision-making concerning ecosystems?
- What tools and methodologies developed and used in the MA can strengthen capacity to assess ecosystems, the services they provide, their impacts on human well-being, and the strengths and weaknesses of response options?

The MA was conducted as a multiscale assessment, with interlinked assessments undertaken at local, watershed, national, regional, and global scales. A global ecosystem assessment cannot easily meet all the needs of decision-makers at national and sub-national scales because the management of any particular ecosystem must be tailored to the particular characteristics of that ecosystem and to the demands placed on it. However, an assessment focused only on a particular ecosystem or particular nation is insufficient because some processes are global and because local goods, services, matter, and energy are often transferred across regions. Each of the component assessments was guided by the MA conceptual framework and benefited from the presence of assessments undertaken at larger and smaller scales. The sub-global assessments were not intended to serve as representative samples of all ecosystems; rather, they were to meet the needs of decision-makers at the scales at which they were undertaken.

The work of the MA was conducted through four working groups, each of which prepared a report of its findings. At the global scale, the Condition and Trends Working Group assessed the state of knowledge on ecosystems, drivers of ecosystem change, ecosystem services, and associated human well-being around the year 2000. The assessment aimed to be comprehensive with regard to ecosystem services, but its coverage is not exhaustive. The Scenarios Working Group considered the possible evolution of ecosystem services during the twenty-first century by developing four global scenarios exploring plausible future changes in drivers, ecosystems, ecosystem services, and human well-being. The Responses Working Group examined the strengths and weaknesses of various response options that have been used to manage ecosystem services and identified promising opportunities for improving human well-being while conserving ecosystems. The report of the Sub-global Assessments Working Group contains lessons learned from the MA sub-global assessments. The first product of the MA—*Ecosystems and Human Well-being: A Framework for Assessment*, published in 2003—outlined the focus, conceptual basis, and methods used in the MA.

Approximately 1,360 experts from 95 countries were involved as authors of the assessment reports, as participants in the sub-global assessments, or as members of the Board of Review Editors. (See Appendix C for the list of coordinating lead authors, sub-global assessment coordinators, and review editors.) The latter group, which involved 80 experts, oversaw the scientific review of the MA reports by governments and experts and ensured that all review comments were appropriately addressed by the authors. All MA findings underwent two rounds of expert and governmental review. Review comments were received from approximately 850 individuals (of which roughly 250 were submitted by authors of other chapters in the MA), although in a number of cases (particularly in the case of governments and MA-affiliated scientific organizations), people submitted collated comments that had been prepared by a number of reviewers in their governments or institutions.

The MA was guided by a Board that included representatives of five international conventions, five U.N. agencies, international scientific organizations, governments, and leaders from the private sector, nongovernmental organizations, and indigenous groups. A 15-member Assessment Panel of leading social and natural scientists oversaw the technical work of the assessment, supported by a secretariat with offices in Europe, North America, South America, Asia, and Africa and coordinated by the United Nations Environment Programme.

The MA is intended to be used:

- to identify priorities for action;
- as a benchmark for future assessments;
- as a framework and source of tools for assessment, planning, and management;
- to gain foresight concerning the consequences of decisions affecting ecosystems;
- to identify response options to achieve human development and sustainability goals;
- to help build individual and institutional capacity to undertake integrated ecosystem assessments and act on the findings; and
- to guide future research.

Because of the broad scope of the MA and the complexity of the interactions between social and natural systems, it proved to be difficult to provide definitive information for some of the issues addressed in the MA. Relatively few ecosystem services have been the focus of research and monitoring and, as a consequence, research findings and data are often inadequate for a detailed global assessment. Moreover, the data and information that are available are generally related to either the characteristics of the ecological system or the characteristics of the social system, not to the all-important interactions between these systems. Finally, the scientific and assessment tools and models available to undertake a cross-scale integrated assessment and to project future changes in ecosystem services are only now being developed. Despite these challenges, the MA was able to provide considerable information relevant to most of the focal questions. And by identifying gaps in data and information that prevent policy-relevant questions from being answered, the assessment can help to guide research and monitoring that may allow those questions to be answered in future assessments.

READER'S GUIDE

This report presents a synthesis and integration of the findings of the four MA Working Groups along with more detailed findings for selected ecosystem services concerning condition and trends and scenarios (see Appendix A) and response options (see Appendix B). Five additional synthesis reports were prepared for ease of use by specific audiences: CBD (biodiversity), UNCCD (desertification), Ramsar Convention (wetlands), business, and the health sector. Each MA sub-global assessment will also produce additional reports to meet the needs of its own audience. The full technical assessment reports of the four MA Working Groups will be published in mid-2005 by Island Press. All printed materials of the assessment, along with core data and a glossary of terminology used in the technical reports, will be available on the Internet at www.MAweb.org. Appendix D lists the acronyms and abbreviations used in this report and includes additional information on sources for some of the Figures. Throughout this report, dollar signs indicate U.S. dollars and tons mean metric tons.

References that appear in parentheses in the body of this synthesis report are to the underlying chapters in the full technical assessment reports of each Working Group. (A list of the assessment report chapters is provided in Appendix E.) To assist the reader, citations to the technical volumes generally specify sections of chapters or specific Boxes, Tables, or Figures, based on final drafts of the chapter. Some chapter subsection numbers may change during final copyediting, however, after this synthesis report has been printed. Bracketed references within the Summary for Decision-makers are to the key questions of this full synthesis report, where additional information on each topic can be found.

In this report, the following words have been used where appropriate to indicate judgmental estimates of certainty, based on the collective judgment of the authors, using the observational evidence, modeling results, and theory that they have examined: very certain (98% or greater probability), high certainty (85–98% probability), medium certainty (65–85% probability), low certainty (52–65% probability), and very uncertain (50–52% probability). In other instances, a qualitative scale to gauge the level of scientific understanding is used: well established, established but incomplete, competing explanations, and speculative. Each time these terms are used they appear in italics.

SUMMARY FOR DECISION-MAKERS



Everyone in the world depends completely on Earth's ecosystems and the services they provide, such as food, water, disease management, climate regulation, spiritual fulfillment, and aesthetic enjoyment. Over the past 50 years, humans have changed these ecosystems more rapidly and extensively than in any comparable period of time in human history, largely to meet rapidly growing demands for food, fresh water, timber, fiber, and fuel. This transformation of the planet has contributed to substantial net gains in human well-being and economic development. But not all regions and groups of people have benefited from this process—in fact, many have been harmed. Moreover, the full costs associated with these gains are only now becoming apparent.

Three major problems associated with our management of the world's ecosystems are already causing significant harm to some people, particularly the poor, and unless addressed will substantially diminish the long-term benefits we obtain from ecosystems:

- First, approximately 60% (15 out of 24) of the ecosystem services examined during the Millennium Ecosystem Assessment are being degraded or used unsustainably, including fresh water, capture fisheries, air and water purification, and the regulation of regional and local climate, natural hazards, and pests. The full costs of the loss and degradation of these ecosystem services are difficult to measure, but the available evidence demonstrates that they are substantial and growing. Many ecosystem services have been degraded as a consequence of actions taken to increase the supply of other services, such as food. These trade-offs often shift the costs of degradation from one group of people to another or defer costs to future generations.

- Second, there is *established but incomplete* evidence that changes being made in ecosystems are increasing the likelihood of nonlinear changes in ecosystems (including accelerating, abrupt, and potentially irreversible changes) that have important consequences for human well-being. Examples of such changes include disease emergence, abrupt alterations in water quality, the creation of “dead zones” in coastal waters, the collapse of fisheries, and shifts in regional climate.

Four Main Findings

- Over the past 50 years, humans have changed ecosystems more rapidly and extensively than in any comparable period of time in human history, largely to meet rapidly growing demands for food, fresh water, timber, fiber, and fuel. This has resulted in a substantial and largely irreversible loss in the diversity of life on Earth.
- The changes that have been made to ecosystems have contributed to substantial net gains in human well-being and economic development, but these gains have been achieved at growing costs in the form of the degradation of many ecosystem services, increased risks of nonlinear changes, and the exacerbation of poverty for some groups of people. These problems, unless addressed, will substantially diminish the benefits that future generations obtain from ecosystems.
- The degradation of ecosystem services could grow significantly worse during the first half of this century and is a barrier to achieving the Millennium Development Goals.
- The challenge of reversing the degradation of ecosystems while meeting increasing demands for their services can be partially met under some scenarios that the MA has considered, but these involve significant changes in policies, institutions, and practices that are not currently under way. Many options exist to conserve or enhance specific ecosystem services in ways that reduce negative trade-offs or that provide positive synergies with other ecosystem services.

■ Third, the harmful effects of the degradation of ecosystem services (the persistent decrease in the capacity of an ecosystem to deliver services) are being borne disproportionately by the poor, are contributing to growing inequities and disparities across groups of people, and are sometimes the principal factor causing poverty and social conflict. This is not to say that ecosystem changes such as increased food production have not also helped to lift many people out of poverty or hunger, but these changes have harmed other individuals and communities, and their plight has been largely overlooked. In all regions, and particularly in sub-Saharan Africa, the condition and management of ecosystem services is a dominant factor influencing prospects for reducing poverty.

The degradation of ecosystem services is already a significant barrier to achieving the Millennium Development Goals agreed to by the international community in September 2000 and the harmful consequences of this degradation could grow significantly worse in the next 50 years. The consumption of ecosystem services, which is unsustainable in many cases, will continue to grow as a consequence of a likely three- to sixfold increase in global GDP by 2050 even while global population growth is expected to slow and level off in mid-century. Most of the important direct drivers of ecosystem change are unlikely to diminish in the first half of the century and two drivers—climate change and excessive nutrient loading—will become more severe.

Already, many of the regions facing the greatest challenges in achieving the MDGs coincide with those facing significant problems of ecosystem degradation. Rural poor people, a primary target of the MDGs, tend to be most directly reliant on ecosystem services and most vulnerable to changes in those services. More generally, any progress achieved in addressing the MDGs of poverty and hunger eradication, improved health, and environmental sustainability is unlikely to be sustained if most of the ecosystem services on which humanity relies continue to be degraded. In contrast, the sound management of ecosystem services provides cost-effective opportunities for addressing multiple development goals in a synergistic manner.

There is no simple fix to these problems since they arise from the interaction of many recognized challenges, including climate change, biodiversity loss, and land degradation, each of which is complex to address in its own right. Past actions to slow or reverse the degradation of ecosystems have yielded significant benefits, but these improvements have generally not kept pace with growing pressures and demands. Nevertheless, there is tremendous scope for action to reduce the severity of these problems in the coming decades. Indeed, three of four detailed scenarios examined by the MA suggest that significant changes in policies, institutions, and practices can mitigate some but not all of the negative consequences of growing pressures on ecosystems. But the changes required are substantial and are not currently under way.

An effective set of responses to ensure the sustainable management of ecosystems requires substantial changes in institutions and

governance, economic policies and incentives, social and behavior factors, technology, and knowledge. Actions such as the integration of ecosystem management goals in various sectors (such as agriculture, forestry, finance, trade, and health), increased transparency and accountability of government and private-sector performance in ecosystem management, elimination of perverse subsidies, greater use of economic instruments and market-based approaches, empowerment of groups dependent on ecosystem services or affected by their degradation, promotion of technologies enabling increased crop yields without harmful environmental impacts, ecosystem restoration, and the incorporation of nonmarket values of ecosystems and their services in management decisions all could substantially lessen the severity of these problems in the next several decades.

The remainder of this Summary for Decision-makers presents the four major findings of the Millennium Ecosystem Assessment on the problems to be addressed and the actions needed to enhance the conservation and sustainable use of ecosystems.

Finding #1: *Over the past 50 years, humans have changed ecosystems more rapidly and extensively than in any comparable period of time in human history, largely to meet rapidly growing demands for food, fresh water, timber, fiber, and fuel. This has resulted in a substantial and largely irreversible loss in the diversity of life on Earth.*

The structure and functioning of the world's ecosystems changed more rapidly in the second half of the twentieth century than at any time in human history. [1]

■ More land was converted to cropland in the 30 years after 1950 than in the 150 years between 1700 and 1850. Cultivated systems (areas where at least 30% of the landscape is in croplands, shifting cultivation, confined livestock production, or freshwater aquaculture) now cover one quarter of Earth's terrestrial surface. (See Figure 1.) Areas of rapid change in forest land cover and land degradation are shown in Figure 2.

■ Approximately 20% of the world's coral reefs were lost and an additional 20% degraded in the last several decades of the twentieth century, and approximately 35% of mangrove area was lost during this time (in countries for which sufficient data exist, which encompass about half of the area of mangroves).

■ The amount of water impounded behind dams quadrupled since 1960, and three to six times as much water is held in reservoirs as in natural rivers. Water withdrawals from rivers and lakes doubled since 1960; most water use (70% worldwide) is for agriculture.

■ Since 1960, flows of reactive (biologically available) nitrogen in terrestrial ecosystems have doubled, and flows of phosphorus have tripled. More than half of all the synthetic nitrogen fertilizer, which was first manufactured in 1913, ever used on the planet has been used since 1985.

Figure 1. EXTENT OF CULTIVATED SYSTEMS, 2000. Cultivated systems cover 24% of the terrestrial surface.





Figure 2. LOCATIONS REPORTED BY VARIOUS STUDIES AS UNDERGOING HIGH RATES OF LAND COVER CHANGE IN THE PAST FEW DECADES (C.SDM)

In the case of forest cover change, the studies refer to the period 1980–2000 and are based on national statistics, remote sensing, and to a limited degree expert opinion. In the case of land cover change resulting from degradation in drylands (desertification), the period is unspecified but inferred to be within the last half-century, and the major study was entirely based on expert opinion, with associated *low certainty*. Change in cultivated area is not shown. Note that areas showing little current change are often locations that have already undergone major historical change (see Figure 1).



■ Since 1750, the atmospheric concentration of carbon dioxide has increased by about 32% (from about 280 to 376 parts per million in 2003), primarily due to the combustion of fossil fuels and land use changes. Approximately 60% of that increase (60 parts per million) has taken place since 1959.

Humans are fundamentally, and to a significant extent irreversibly, changing the diversity of life on Earth, and most of these changes represent a loss of biodiversity. [1]

■ More than two thirds of the area of 2 of the world's 14 major terrestrial biomes and more than half of the area of 4 other biomes had been converted by 1990, primarily to agriculture. (See Figure 3.)

■ Across a range of taxonomic groups, either the population size or range or both of the majority of species is currently declining.

■ The distribution of species on Earth is becoming more homogenous; in other words, the set of species in any one region of the world is becoming more similar to the set in other regions primarily as a result of introductions of species, both intentionally and inadvertently in association with increased travel and shipping.


■ The number of species on the planet is declining. Over the past few hundred years, humans have increased the species extinction rate by as much as 1,000 times over background rates typical over the planet's history (*medium certainty*). (See Figure 4.) Some 10–30% of mammal, bird, and amphibian species are currently threatened with extinction (*medium to high certainty*). Freshwater ecosystems tend to have the highest proportion of species threatened with extinction.

■ Genetic diversity has declined globally, particularly among cultivated species.

Most changes to ecosystems have been made to meet a dramatic growth in the demand for food, water, timber, fiber, and fuel. [2] Some ecosystem changes have been the inadvertent result of activities unrelated to the use of ecosystem services, such as the construction of roads, ports, and cities and the discharge of pollutants. But most ecosystem changes were the direct or indirect result of changes made to meet growing demands for ecosystem services, and in particular growing demands for food, water, timber, fiber, and fuel (fuelwood and hydropower).

Figure 3. CONVERSION OF TERRESTRIAL BIOMES^a
(Adapted from C4, S10)

It is not possible to estimate accurately the extent of different biomes prior to significant human impact, but it is possible to determine the “potential” area of biomes based on soil and climatic conditions. This Figure shows how much of that potential area is estimated to have been converted by 1950 (*medium certainty*), how much was converted between 1950 and 1990 (*medium certainty*), and how much would be converted under the four MA scenarios (*low certainty*) between 1990 and 2050. Mangroves are not included here because the area was too small to be accurately assessed. Most of the conversion of these biomes is to cultivated systems.



^a A biome is the largest unit of ecological classification that is convenient to recognize below the entire globe, such as temperate broadleaf forests or montane grasslands. A biome is a widely used ecological categorization, and because considerable ecological data have been reported and modeling undertaken using this categorization, some information in this assessment can only be reported based on biomes. Whenever possible, however, the MA reports information using 10 socioecological systems, such as forest, cultivated, coastal, and marine, because these correspond to the regions of responsibility of different government ministries and because they are the categories used within the Convention on Biological Diversity.

^b According to the four MA scenarios. For 2050 projections, the average value of the projections under the four scenarios is plotted and the error bars (black lines) represent the range of values from the different scenarios.

Source: Millennium Ecosystem Assessment

Between 1960 and 2000, the demand for ecosystem services grew significantly as world population doubled to 6 billion people and the global economy increased more than sixfold. To meet this demand, food production increased by roughly two-and-a-half times, water use doubled, wood harvests for pulp and paper production tripled, installed hydropower capacity doubled, and timber production increased by more than half.

The growing demand for these ecosystem services was met both by consuming an increasing fraction of the available supply (for example, diverting more water for irrigation or capturing more fish from the sea) and by raising the production of some services, such as crops and livestock. The latter has been accomplished through the use of new technologies (such as new crop varieties, fertilization, and irrigation) as well as through increasing the area managed for the services in the case of crop and livestock production and aquaculture.

Finding #2: *The changes that have been made to ecosystems have contributed to substantial net gains in human well-being and economic development, but these gains have been achieved at growing costs in the form of the degradation of many ecosystem services, increased risks of nonlinear changes, and the exacerbation of poverty for some groups of people. These problems, unless addressed, will substantially diminish the benefits that future generations obtain from ecosystems.*

In the aggregate, and for most countries, changes made to the world's ecosystems in recent decades have provided substantial benefits for human well-being and national development. [3] Many of the most significant changes to ecosystems have been essential to meet growing needs for food and water; these


Figure 4. SPECIES EXTINCTION RATES (Adapted from C4 Fig 4.22)

“Distant past” refers to average extinction rates as estimated from the fossil record. “Recent past” refers to extinction rates calculated from known extinctions of species (lower estimate) or known extinctions plus “possibly extinct” species (upper bound). A species is considered to be “possibly extinct” if it is believed by experts to be extinct but extensive surveys have not yet been undertaken to confirm its disappearance.

“Future” extinctions are model-derived estimates using a variety of techniques, including species-area models, rates at which species are shifting to increasingly more threatened categories, extinction probabilities associated with the IUCN categories of threat, impacts of projected habitat loss on species currently threatened with habitat loss, and correlation of species loss with energy consumption. The time frame and species groups involved differ among the “future” estimates, but in general refer to either future loss of species based on the level of threat that exists

today or current and future loss of species as a result of habitat changes taking place over the period of roughly 1970 to 2050. Estimates based on the fossil record are *low certainty*; lower-bound estimates for known extinctions are *high certainty* and upper-bound estimates are *medium certainty*; lower-bound estimates for modeled extinctions are *low certainty* and upper-bound estimates are *speculative*. The rate of known extinctions of species in the past century is roughly 50–500 times greater than the extinction rate calculated from the fossil record of 0.1–1 extinctions per 1,000 species per 1,000 years. The rate is up to 1,000 times higher than the background extinction rates if possibly extinct species are included.

Extinctions per thousand species per millennium



Source: Millennium Ecosystem Assessment

changes have helped reduce the proportion of malnourished people and improved human health. Agriculture, including fisheries and forestry, has been the mainstay of strategies for the development of countries for centuries, providing revenues that have enabled investments in industrialization and poverty alleviation. Although the value of food production in 2000 was only about 3% of gross world product, the agricultural labor force accounts for approximately 22% of the world's population, half the world's total labor force, and 24% of GDP in countries with per capita incomes of less than \$765 (the low-income developing countries, as defined by the World Bank).

These gains have been achieved, however, at growing costs in the form of the degradation of many ecosystem services, increased risks of nonlinear changes in ecosystems, the exacerbation of poverty for some people, and growing inequities and disparities across groups of people.

Degradation and Unsustainable Use of Ecosystem Services

Approximately 60% (15 out of 24) of the ecosystem services evaluated in this assessment (including 70% of regulating and cultural services) are being degraded or used unsustainably. [2] (See Table 1.) Ecosystem services that have been degraded over the past 50 years include capture fisheries, water supply, waste treatment and detoxification, water purification, natural hazard protection, regulation of air quality, regulation of regional and local climate, regulation of erosion, spiritual fulfillment, and aesthetic enjoyment. The use of two ecosystem services—capture fisheries and fresh water—is now well beyond levels that can be sustained even at current demands, much less future ones. At least one quarter of important commercial fish stocks are overharvested (*high certainty*). (See Figures 5, 6, and 7.) From 5% to possibly 25% of global freshwater use exceeds long-term accessible supplies and is now met either through engineered water transfers or overdraft of groundwater supplies (*low to medium certainty*). Some 15–35% of irrigation withdrawals exceed supply rates and are therefore unsustainable (*low to medium certainty*). While 15 services have been degraded, only 4 have been enhanced in the past 50 years, three of which involve food production: crops, livestock, and aquaculture. Terrestrial ecosystems were on average a net source of CO₂ emissions during the nineteenth and early twentieth centuries, but became a net sink around the middle of the last century, and thus in the last 50 years the role of ecosystems in regulating global climate through carbon sequestration has also been enhanced.

Actions to increase one ecosystem service often cause the degradation of other services. [2, 6] For example, because actions to increase food production typically involve increased use of water and fertilizers or expansion of the area of cultivated land,

these same actions often degrade other ecosystem services, including reducing the availability of water for other uses, degrading water quality, reducing biodiversity, and decreasing forest cover (which in turn may lead to the loss of forest products and the release of greenhouse gasses). Similarly, the conversion of forest to agriculture can significantly change the frequency and magnitude of floods, although the nature of this impact depends on the characteristics of the local ecosystem and the type of land cover change.

The degradation of ecosystem services often causes significant harm to human well-being. [3, 6] The information available to assess the consequences of changes in ecosystem services for human well-being is relatively limited. Many ecosystem services have not been monitored, and it is also difficult to estimate the influence of changes in ecosystem services relative to other social, cultural, and economic factors that also affect human well-being. Nevertheless, the following types of evidence demonstrate that the harmful effects of the degradation of ecosystem services on livelihoods, health, and local and national economies are substantial.

■ *Most resource management decisions are most strongly influenced by ecosystem services entering markets; as a result, the nonmarketed benefits are often lost or degraded. These nonmarketed benefits are often high and sometimes more valuable than the marketed ones.* For example, one of the most comprehensive studies to date, which examined the marketed and nonmarketed economic values associated with forests in eight Mediterranean countries, found that timber and fuelwood generally accounted for less than a third of total economic value of forests in each country. (See Figure 8.) Values associated with non-wood forest products, recreation, hunting, watershed protection, carbon sequestration, and passive use (values independent of direct uses) accounted for between 25% and 96% of the total economic value of the forests.

■ *The total economic value associated with managing ecosystems more sustainably is often higher than the value associated with the conversion of the ecosystem through farming, clear-cut logging, or other intensive uses.* Relatively few studies have compared the total economic value (including values of both marketed and nonmarketed ecosystem services) of ecosystems under alternate management regimes, but some of the studies that do exist have found that the benefit of managing the ecosystem more sustainably exceeded that of converting the ecosystem. (See Figure 9.)

■ *The economic and public health costs associated with damage to ecosystem services can be substantial.*

- The early 1990s collapse of the Newfoundland cod fishery due to overfishing resulted in the loss of tens of thousands of jobs and cost at least \$2 billion in income support and retraining.
- In 1996, the cost of U.K. agriculture resulting from the damage that agricultural practices cause to water (pollution and eutrophication, a process whereby excessive plant growth depletes oxygen in the water), air (emissions of greenhouse gases), soil (off-site erosion damage, emissions

Table 1. GLOBAL STATUS OF PROVISIONING, REGULATING, AND CULTURAL ECOSYSTEM SERVICES EVALUATED IN THE MA

Status indicates whether the condition of the service globally has been enhanced (if the productive capacity of the service has been increased, for example) or degraded in the recent past. Definitions of “enhanced” and “degraded” are provided in the note below. A fourth category, supporting services, is not included here as they are not used directly by people.

Service	Sub-category	Status	Notes
Provisioning Services			
Food	crops	▲	substantial production increase
	livestock	▲	substantial production increase
	capture fisheries	▼	declining production due to overharvest
	aquaculture	▲	substantial production increase
	wild foods	▼	declining production
Fiber	timber	+/-	forest loss in some regions, growth in others
	cotton, hemp, silk	+/-	declining production of some fibers, growth in others
	wood fuel	▼	declining production
Genetic resources		▼	lost through extinction and crop genetic resource loss
Biochemicals, natural medicines, pharmaceuticals		▼	lost through extinction, overharvest
Fresh water		▼	unsustainable use for drinking, industry, and irrigation; amount of hydro energy unchanged, but dams increase ability to use that energy
Regulating Services			
Air quality regulation		▼	decline in ability of atmosphere to cleanse itself
Climate regulation	global	▲	net source of carbon sequestration since mid-century
	regional and local	▼	preponderance of negative impacts
Water regulation		+/-	varies depending on ecosystem change and location
Erosion regulation		▼	increased soil degradation
Water purification and waste treatment		▼	declining water quality
Disease regulation		+/-	varies depending on ecosystem change
Pest regulation		▼	natural control degraded through pesticide use
Pollination		▼ ^a	apparent global decline in abundance of pollinators
Natural hazard regulation		▼	loss of natural buffers (wetlands, mangroves)
Cultural Services			
Spiritual and religious values		▼	rapid decline in sacred groves and species
Aesthetic values		▼	decline in quantity and quality of natural lands
Recreation and ecotourism		+/-	more areas accessible but many degraded

Note: For provisioning services, we define enhancement to mean increased production of the service through changes in area over which the service is provided (e.g., spread of agriculture) or increased production per unit area. We judge the production to be degraded if the current use exceeds sustainable levels. For regulating and supporting services, enhancement refers to a change in the service that leads to greater benefits for people (e.g., the service of disease regulation could be improved by eradication of a vector known to transmit a disease to people). Degradation of regulating and supporting services means a reduction in the benefits obtained from the service, either through a change in the service (e.g., mangrove loss reducing the storm protection benefits of an ecosystem) or through human pressures on the service exceeding its limits (e.g., excessive pollution exceeding the capability of ecosystems to maintain water quality). For cultural services, enhancement refers to a change in the ecosystem features that increase the cultural (recreational, aesthetic, spiritual, etc.) benefits provided by the ecosystem.

^a Indicates low to medium certainty. All other trends are medium to high certainty.

Figure 5. ESTIMATED GLOBAL MARINE FISH CATCH, 1950–2001 (C18 Fig 18.3)

In this Figure, the catch reported by governments is in some cases adjusted to correct for likely errors in data.




Figure 7. TREND IN MEAN DEPTH OF CATCH SINCE 1950. FISHERIES CATCHES INCREASINGLY ORIGINATE FROM DEEP AREAS (Data from C18 Fig 18.5)





Figure 6. DECLINE IN TROPIC LEVEL OF FISHERIES CATCH SINCE 1950 (C18)

A trophic level of an organism is its position in a food chain. Levels are numbered according to how far particular organisms are along the chain from the primary producers at level 1, to herbivores (level 2), to predators (level 3), to carnivores or top carnivores (level 4 or 5). Fish at higher trophic levels are typically of higher economic value. The decline in the trophic level harvested is largely a result of the overharvest of fish at higher trophic levels.



of greenhouse gases), and biodiversity was \$2.6 billion, or 9% of average yearly gross farm receipts for the 1990s. Similarly, the damage costs of freshwater eutrophication alone in England and Wales (involving factors including reduced value of waterfront dwellings, water treatment costs, reduced recreational value of water bodies, and tourism losses) was estimated to be \$105–160 million per year in the 1990s, with an additional \$77 million a year being spent to address those damages.

- The incidence of diseases of marine organisms and the emergence of new pathogens is increasing, and some of these, such as ciguatera, harm human health. Episodes of harmful (including toxic) algal blooms in coastal waters are increasing in frequency and intensity, harming other marine resources such as fisheries as well as human health. In a particularly severe outbreak in Italy in 1989, harmful algal blooms cost the coastal aquaculture industry \$10 million and the Italian tourism industry \$11.4 million.
- The frequency and impact of floods and fires has increased significantly in the past 50 years, in part due to ecosystem changes. Examples are the increased susceptibility of coastal populations to tropical storms when mangrove forests are cleared and the increase in downstream flooding that followed land use changes in the upper Yangtze River. Annual economic losses from extreme events increased tenfold from the 1950s to approximately \$70 billion in 2003, of which natural catastrophes (floods, fires, storms, drought, earthquakes) accounted for 84% of insured losses.


■ *The impact of the loss of cultural services is particularly difficult to measure, but it is especially important for many people.* Human cultures, knowledge systems, religions, and social interactions have been strongly influenced by ecosystems. A number of the MA sub-global assessments found that spiritual and cultural values of ecosystems were as important as other services for many local communities, both in developing countries (the importance of sacred groves of forest in India, for example) and industrial ones (the importance of urban parks, for instance).

The degradation of ecosystem services represents loss of a capital asset. [3] Both renewable resources such as ecosystem services and nonrenewable resources such as mineral deposits, some soil nutrients, and fossil fuels are capital assets. Yet traditional national accounts do not include measures of resource depletion or of the degradation of these resources. As a result, a country could cut its forests and deplete its fisheries, and this would show only as a positive gain in GDP (a measure of current economic well-being) without registering the corresponding decline in assets (wealth) that is the more appropriate measure of future economic well-being. Moreover, many ecosystem services (such as fresh water in aquifers and the use of the atmosphere as a sink for pollutants) are available freely to those who use them, and so again their degradation is not reflected in standard economic measures.

When estimates of the economic losses associated with the depletion of natural assets are factored into measurements of the total wealth of nations, they significantly change the balance

Figure 8. ANNUAL FLOW OF BENEFITS FROM FORESTS IN SELECTED COUNTRIES
(Adapted from C5 Box 5.2)


In most countries, the marketed values of ecosystems associated with timber and fuelwood production are less than one third of the total economic value, including nonmarketed values such as carbon sequestration, watershed protection, and recreation.



sheet of countries with economies significantly dependent on natural resources. For example, countries such as Ecuador, Ethiopia, Kazakhstan, Democratic Republic of Congo, Trinidad and Tobago, Uzbekistan, and Venezuela that had positive growth in net savings in 2001, reflecting a growth in the net wealth of the country, actually experienced a loss in net savings when depletion of natural resources (energy and forests) and estimated damages from carbon emissions (associated with contributions to climate change) were factored into the accounts.

Figure 9. ECONOMIC BENEFITS UNDER ALTERNATE MANAGEMENT PRACTICES (C5 Box 5.2)

In each case, the net benefits from the more sustainably managed ecosystem are greater than those from the converted ecosystem, even though the private (market) benefits would be greater from the converted ecosystem. (Where ranges of values are given in the original source, lower estimates are plotted here.)



Source: Millennium Ecosystem Assessment

While degradation of some services may sometimes be warranted to produce a greater gain in other services, often more degradation of ecosystem services takes place than is in society's interests because many of the services degraded are "public goods." [3] Although people benefit from ecosystem services such as the regulation of air and water quality or the presence of an

aesthetically pleasing landscape, there is no market for these services and no one person has an incentive to pay to maintain the good. And when an action results in the degradation of a service that harms other individuals, no market mechanism exists (nor, in many cases, could it exist) to ensure that the individuals harmed are compensated for the damages they suffer.

Wealthy populations cannot be insulated from the degradation of ecosystem services. [3] Agriculture, fisheries, and forestry once formed the bulk of national economies, and the control of natural resources dominated policy agendas. But while these natural resource industries are often still important, the relative economic and political significance of other industries in industrial countries has grown over the past century as a result of the ongoing transition from agricultural to industrial and service economies, urbanization, and the development of new technologies to increase the production of some services and provide substitutes for others. Nevertheless, the degradation of ecosystem services influences human well-being in industrial regions and among wealthy populations in developing countries in many ways:

- The physical, economic, or social impacts of ecosystem service degradation may cross boundaries. (See Figure 10.) For example, land degradation and associated dust storms or fires in one country can degrade air quality in other countries nearby.
- Degradation of ecosystem services exacerbates poverty in developing countries, which can affect neighboring industrial countries by slowing regional economic growth and contributing to the outbreak of conflicts or the migration of refugees.
- Changes in ecosystems that contribute to greenhouse gas emissions contribute to global climate changes that affect all countries.
- Many industries still depend directly on ecosystem services. The collapse of fisheries, for example, has harmed many communities in industrial countries. Prospects for the forest, agriculture, fishing, and ecotourism industries are all directly tied to ecosystem services, while other sectors such as insurance, banking, and health are strongly, if less directly, influenced by changes in ecosystem services.

■ Wealthy populations of people are insulated from the harmful effects of some aspects of ecosystem degradation, but not all. For example, substitutes are typically not available when cultural services are lost.

■ Even though the relative economic importance of agriculture, fisheries, and forestry is declining in industrial countries, the importance of other ecosystem services such as aesthetic enjoyment and recreational options is growing.

It is difficult to assess the implications of ecosystem changes and to manage ecosystems effectively because many of the effects are slow to become apparent, because they may be expressed primarily at some distance from where the ecosystem was changed, and because the costs and benefits of changes often accrue to different sets of stakeholders. [7] Substantial inertia (delay in the response of a system to a disturbance) exists in ecological systems. As a result, long time lags often occur between a change in a driver and the time when the full consequences of that change become apparent. For example, phosphorus is accumulating in large quantities in many agricultural soils, threatening rivers, lakes, and coastal oceans with increased eutrophication. But it may take years or decades for the full impact of the phosphorus to become apparent through erosion and other processes. Similarly, it will take centuries for global temperatures to reach equilibrium with changed concentrations of greenhouse gases in the atmosphere and even more time for biological systems to respond to the changes in climate.

Moreover, some of the impacts of ecosystem changes may be experienced only at some distance from where the change occurred. For example, changes in upstream catchments affect water flow and water quality in downstream regions; similarly, the loss of an important fish nursery area in a coastal wetland may diminish fish catch some distance away. Both the inertia in ecological systems and the temporal and spatial separation of costs and benefits of ecosystem changes often result in situations where the individuals experiencing harm from ecosystem changes (future generations, say, or downstream landowners) are not the same as the individuals gaining the benefits. These temporal and spatial patterns make it extremely difficult to fully assess costs and benefits associated with ecosystem changes or to attribute costs and benefits to different stakeholders. Moreover, the institutional arrangements now in place to manage ecosystems are poorly designed to cope with these challenges.

Increased Likelihood of Nonlinear (Stepped) and Potentially Abrupt Changes in Ecosystems


There is *established but incomplete* evidence that changes being made in ecosystems are increasing the likelihood of nonlinear changes in ecosystems (including accelerating, abrupt, and potentially irreversible changes), with important consequences for human well-being. [7] Changes in ecosystems generally take place gradually. Some changes are nonlinear, however: once a threshold is crossed, the system changes to a very different state. And these nonlinear changes are sometimes abrupt; they can also be large in magnitude and difficult, expensive, or impossible to reverse. Capabilities for predicting some nonlinear changes are improving, but for most ecosystems and for most potential nonlinear changes, while science can often warn of increased risks of change it cannot predict the thresholds at which the change will be encountered. Examples of large-magnitude nonlinear changes include:

■ *Disease emergence.* If, on average, each infected person infects at least one other person, then an epidemic spreads, while if the infection is transferred on average to less than one person, the epidemic dies out. During the 1997–98 El Niño, excessive flooding caused cholera epidemics in Djibouti, Somalia, Kenya, Tanzania, and Mozambique. Warming of the African Great Lakes due to climate change may create conditions that increase the risk of cholera transmission in the surrounding countries.

■ *Eutrophication and hypoxia.* Once a threshold of nutrient loading is achieved, changes in freshwater and coastal ecosystems can be abrupt and extensive, creating harmful algal blooms (including blooms of toxic species) and sometimes leading to the formation of oxygen-depleted zones, killing most animal life.

Figure 10. DUST CLOUD OFF THE NORTHWEST COAST OF AFRICA, MARCH 6, 2004

In this image, the storm covers about one fifth of Earth's circumference. The dust clouds travel thousands of kilometers and fertilize the water off the west coast of Florida with iron. This has been linked to blooms of toxic algae in the region and respiratory problems in North America and has affected coral reefs in the Caribbean. Degradation of drylands exacerbates problems associated with dust storms.



Source: National Aeronautics and Space Administration, Earth Observatory

■ **Fisheries collapse.** For example, the Atlantic cod stocks off the east coast of Newfoundland collapsed in 1992, forcing the closure of the fishery after hundreds of years of exploitation. (See Figure 11.) Most important, depleted stocks may take years to recover, or not recover at all, even if harvesting is significantly reduced or eliminated entirely.

■ **Species introductions and losses.** The introduction of the zebra mussel into aquatic systems in the United States, for instance, resulted in the extirpation of native clams in Lake St. Clair and annual costs of \$100 million to the power industry and other users.

■ **Regional climate change.** Deforestation generally leads to decreased rainfall. Since forest existence crucially depends on rainfall, the relationship between forest loss and precipitation decrease can form a positive feedback, which, under certain conditions, can lead to a nonlinear change in forest cover.

The growing bushmeat trade poses particularly significant threats associated with nonlinear changes, in this case accelerating rates of change. [7] Growth in the use and trade of bushmeat is placing increasing pressure on many species, especially in Africa and Asia. While the population size of harvested species may decline gradually with increasing harvest for some time, once the harvest exceeds sustainable levels, the rate of decline of populations of the harvested species will tend to accelerate. This could place them at risk of extinction and also reduce the food supply of people dependent on these resources in the longer term. At the same time, the bushmeat trade involves relatively high levels of interaction between humans and some relatively closely related wild animals that are eaten. Again, this increases the risk of a nonlinear change, in this case the emergence of new and serious pathogens. Given the speed and magnitude of international travel today, new pathogens could spread rapidly around the world.

The increased likelihood of these nonlinear changes stems from the loss of biodiversity and growing pressures from multiple direct drivers of ecosystem change. [7] The loss of species and genetic diversity decreases the resilience of ecosystems, which is the level of disturbance that an ecosystem can undergo without crossing a threshold to a different

structure or functioning. In addition, growing pressures from drivers such as overharvesting, climate change, invasive species, and nutrient loading push ecosystems toward thresholds that they might otherwise not encounter.


Exacerbation of Poverty for Some Individuals and Groups of People and Contribution to Growing Inequities and Disparities across Groups of People

Despite the progress achieved in increasing the production and use of some ecosystem services, levels of poverty remain high, inequities are growing, and many people still do not have a sufficient supply of or access to ecosystem services. [3]

■ In 2001, 1.1 billion people survived on less than \$1 per day of income, with roughly 70% of them in rural areas where they are highly dependent on agriculture, grazing, and hunting for subsistence.

Figure 11. COLLAPSE OF ATLANTIC COD STOCKS OFF THE EAST COAST OF NEWFOUNDLAND IN 1992 (CF Box 2.4)

This collapse forced the closure of the fishery after hundreds of years of exploitation. Until the late 1950s, the fishery was exploited by migratory seasonal fleets and resident inshore small-scale fishers. From the late 1950s, offshore bottom trawlers began exploiting the deeper part of the stock, leading to a large catch increase and a strong decline in the underlying biomass. Internationally agreed quotas in the early 1970s and, following the declaration by Canada of an Exclusive Fishing Zone in 1977, national quota systems ultimately failed to arrest and reverse the decline. The stock collapsed to extremely low levels in the late 1980s and early 1990s, and a moratorium on commercial fishing was declared in June 1992. A small commercial inshore fishery was reintroduced in 1998, but catch rates declined and the fishery was closed indefinitely in 2003.



Source: Millennium Ecosystem Assessment

■ Inequality in income and other measures of human well-being has increased over the past decade. A child born in sub-Saharan Africa is 20 times more likely to die before age 5 than a child born in an industrial country, and this disparity is higher than it was a decade ago. During the 1990s, 21 countries experienced declines in their rankings in the Human Development Index (an aggregate measure of economic well-being, health, and education); 14 of them were in sub-Saharan Africa.

■ Despite the growth in per capita food production in the past four decades, an estimated 852 million people were undernourished in 2000–02, up 37 million from the period 1997–99. South Asia and sub-Saharan Africa, the regions with the largest numbers of undernourished people, are also the regions where growth in per capita food production has lagged the most. Most notably, per capita food production has declined in sub-Saharan Africa.

■ Some 1.1 billion people still lack access to improved water supply, and more than 2.6 billion lack access to improved sanitation. Water scarcity affects roughly 1–2 billion people worldwide. Since 1960, the ratio of water use to accessible supply has grown by 20% per decade.

The degradation of ecosystem services is harming many of the world's poorest people and is sometimes the principal factor causing poverty. [3, 6]

■ Half the urban population in Africa, Asia, Latin America, and the Caribbean suffers from one or more diseases associated with inadequate water and sanitation. Worldwide, approximately 1.7 million people die annually as a result of inadequate water, sanitation, and hygiene.

■ The declining state of capture fisheries is reducing an inexpensive source of protein in developing countries. Per capita fish consumption in developing countries, excluding China, declined between 1985 and 1997.

■ Desertification affects the livelihoods of millions of people, including a large portion of the poor in drylands.

The pattern of “winners” and “losers” associated with ecosystem changes—and in particular the impact of ecosystem changes on poor people, women, and indigenous peoples—has not been adequately taken into account in management decisions. [3, 6] Changes in ecosystems typically yield benefits for some people and exact costs on others who may either lose access to resources or livelihoods or be affected by externalities associated with the change. For several reasons, groups such as the poor, women, and indigenous communities have tended to be harmed by these changes.

■ Many changes in ecosystem management have involved the privatization of what were formerly common pool resources. Individuals who depended on those resources (such as indigenous peoples, forest-dependent communities, and other groups relatively marginalized from political and economic sources of power) have often lost rights to the resources.

■ Some of the people and places affected by changes in ecosystems and ecosystem services are highly vulnerable and poorly equipped to cope with the major changes in ecosystems that may occur. Highly vulnerable groups include those whose needs for

ecosystem services already exceed the supply, such as people lacking adequate clean water supplies, and people living in areas with declining per capita agricultural production.

■ Significant differences between the roles and rights of men and women in many societies lead to increased vulnerability of women to changes in ecosystem services.


■ The reliance of the rural poor on ecosystem services is rarely measured and thus typically overlooked in national statistics and poverty assessments, resulting in inappropriate strategies that do not take into account the role of the environment in poverty reduction. For example, a recent study that synthesized data from 17 countries found that 22% of household income for rural communities in forested regions comes from sources typically not included in national statistics, such as harvesting wild food, fuelwood, fodder, medicinal plants, and timber. These activities generated a much higher proportion of poorer families' total income than of wealthy families', and this income was of particular significance in periods of both predictable and unpredictable shortfalls in other livelihood sources.

Development prospects in dryland regions of developing countries are especially dependent on actions to avoid the degradation of ecosystems and slow or reverse degradation where it is occurring. [3, 5] Dryland systems cover about 41% of Earth's land surface and more than 2 billion people inhabit them, more than 90% of whom are in developing countries. Dryland ecosystems (encompassing both rural and urban regions of drylands) experienced the highest population growth rate in the 1990s of any of the systems examined in the MA. (See Figure 12.) Although drylands are home to about one third of the human population, they have only 8% of the world's renewable water supply. Given the low and variable rainfall, high temperatures, low soil organic matter, high costs of delivering services such as electricity or piped water, and limited investment in infrastructure due to the low population density, people living in drylands face many challenges. They also tend to have the lowest levels of human well-being, including the lowest per capita GDP and the highest infant mortality rates.

The combination of high variability in environmental conditions and relatively high levels of poverty leads to situations where people can be highly vulnerable to changes in ecosystems, although the presence of these conditions has led to the development of very resilient land management strategies. Pressures on dryland ecosystems already exceed sustainable levels for some ecosystem services, such as soil formation and water supply, and are growing. Per capita water availability is currently only two thirds of the level required for minimum levels of human well-being. Approximately 10–20% of the world's drylands are degraded (*medium certainty*) directly harming the people living in these areas and indirectly harming a larger population through biophysical impacts (dust storms, greenhouse gas emissions, and regional climate change) and through socioeconomic impacts

Figure 12. HUMAN POPULATION GROWTH RATES, 1990–2000, AND PER CAPITA GDP AND BIOLOGICAL PRODUCTIVITY IN 2000 IN MA ECOLOGICAL SYSTEMS (C.SDM)

MA systems with the lowest net primary productivity and lowest GDP tended to have the highest population growth rates between 1990 and 2000. Urban, inland water, and marine systems are not included due to the somewhat arbitrary nature of determining net primary productivity of the system (urban) or population growth and GDP (freshwater and marine) for them.



Sources: Millennium Ecosystem Assessment

(human migration and deepening poverty sometimes contributing to conflict and instability). Despite these tremendous challenges, people living in drylands and their land management systems have a proven resilience and the capability of preventing land degradation, although this can be either undermined or enhanced by public policies and development strategies.

Finding #3: *The degradation of ecosystem services could grow significantly worse during the first half of this century and is a barrier to achieving the Millennium Development Goals.*

The MA developed four scenarios to explore plausible futures for ecosystems and human well-being. (See Box 1.) The scenarios explored two global development paths, one in which the world becomes increasingly globalized and the other in which it becomes increasingly regionalized, as well as two different approaches to ecosystem management, one in which actions are reactive and most problems are addressed only after they become obvious and the other in which ecosystem management is proactive and policies deliberately seek to maintain ecosystem services for the long term.

Most of the direct drivers of change in ecosystems currently remain constant or are growing in intensity in most ecosystems. (See Figure 13.) In all four MA scenarios, the pressures on ecosystems are projected to continue to grow during the first half of this century. [4, 5] The most important direct drivers of change in ecosystems are habitat change (land use change and physical modification of rivers or water withdrawal from rivers), overexploitation, invasive alien species, pollution, and climate change. These direct drivers are often synergistic. For example, in some locations land use change can result in greater nutrient loading (if the land is converted to high-intensity agriculture), increased emissions of greenhouse gases (if forest is cleared), and increased numbers of invasive species (due to the disturbed habitat).

■ **Habitat transformation, particularly from conversion to agriculture:** Under the MA scenarios, a further 10–20% of grassland and forestland is projected to be converted between 2000 and 2050 (primarily to agriculture), as Figure 2 illustrated. The projected land conversion is concentrated in low-income countries and dryland regions. Forest cover is projected to continue to increase within industrial countries.

■ *Overexploitation, especially overfishing:* Over much of the world, the biomass of fish targeted in fisheries (including that of both the target species and those caught incidentally) has been reduced by 90% relative to levels prior to the onset of industrial fishing, and the fish being harvested are increasingly coming from the less valuable lower trophic levels as populations of higher trophic level species are depleted, as shown in Figure 6. These pressures continue to grow in all the MA scenarios.

■ *Invasive alien species:* The spread of invasive alien species and disease organisms continues to increase because of both deliberate translocations and accidental introductions related to growing trade and travel, with significant harmful consequences to native species and many ecosystem services.

■ *Pollution, particularly nutrient loading:* Humans have already doubled the flow of reactive nitrogen on the continents, and some projections suggest that this may increase by roughly a

further two thirds by 2050. (See Figure 14.) Three out of four MA scenarios project that the global flux of nitrogen to coastal ecosystems will increase by a further 10–20% by 2030 (*medium certainty*), with almost all of this increase occurring in developing countries. Excessive flows of nitrogen contribute to eutrophication of freshwater and coastal marine ecosystems and acidification of freshwater and terrestrial ecosystems (with implications for biodiversity in these ecosystems). To some degree, nitrogen also plays a role in creation of ground-level ozone (which leads to loss of agricultural and forest productivity), destruction of ozone in the stratosphere (which leads to depletion of the ozone layer and increased UV-B radiation on Earth, causing increased incidence of skin cancer), and climate change. The resulting health effects include the consequences of ozone pollution on asthma and respiratory function, increased allergies and asthma due to increased pollen production, the risk of blue-baby syndrome,

Box 1. MA SCENARIOS

The MA developed four scenarios to explore plausible futures for ecosystems and human well-being based on different assumptions about driving forces of change and their possible interactions:

Global Orchestration – This scenario depicts a globally connected society that focuses on global trade and economic liberalization and takes a reactive approach to ecosystem problems but that also takes strong steps to reduce poverty and inequality and to invest in public goods such as infrastructure and education. Economic growth in this scenario is the highest of the four scenarios, while it is assumed to have the lowest population in 2050.

Order from Strength – This scenario represents a regionalized and fragmented world, concerned with security and protection, emphasizing primarily regional markets, paying little attention to public goods, and taking a reactive approach to ecosystem problems. Economic growth rates are the lowest of the scenarios (particularly low in developing countries) and decrease with time, while population growth is the highest.

Adapting Mosaic – In this scenario, regional watershed-scale ecosystems are the focus of political and economic activity. Local institutions are strengthened and local ecosystem management strategies are common; societies develop a strongly proactive approach to the management of ecosystems. Economic growth rates are somewhat low initially but

increase with time, and population in 2050 is nearly as high as in *Order from Strength*.

TechnoGarden – This scenario depicts a globally connected world relying strongly on environmentally sound technology, using highly managed, often engineered, ecosystems to deliver ecosystem services, and taking a proactive approach to the management of ecosystems in an effort to avoid problems. Economic growth is relatively high and accelerates, while population in 2050 is in the mid-range of the scenarios.

The scenarios are not predictions; instead they were developed to explore the unpredictable features of change in drivers and ecosystem services. No scenario represents business as usual, although all begin from current conditions and trends.

Both quantitative models and qualitative analyses were used to develop the scenarios. For some drivers (such as land use change and carbon emissions) and ecosystem services (water withdrawals, food production), quantitative projections were calculated using established, peer-reviewed global models. Other drivers (such as rates of technological change and economic growth), ecosystem services (particularly supporting and cultural services, such as soil formation and recreational opportunities), and human well-being indicators (such as human health and social relations) were estimated qualitatively. In general, the quantitative models used for these scenarios addressed incremen-

tal changes but failed to address thresholds, risk of extreme events, or impacts of large, extremely costly, or irreversible changes in ecosystem services. These phenomena were addressed qualitatively by considering the risks and impacts of large but unpredictable ecosystem changes in each scenario.

Three of the scenarios – *Global Orchestration*, *Adapting Mosaic*, and *TechnoGarden* incorporate significant changes in policies aimed at addressing sustainable development challenges. In *Global Orchestration* trade barriers are eliminated, distorting subsidies are removed, and a major emphasis is placed on eliminating poverty and hunger. In *Adapting Mosaic*, by 2010, most countries are spending close to 13% of their GDP on education (as compared to an average of 3.5% in 2000), and institutional arrangements to promote transfer of skills and knowledge among regional groups proliferate. In *TechnoGarden* policies are put in place to provide payment to individuals and companies that provide or maintain the provision of ecosystem services. For example, in this scenario, by 2015, roughly 50% of European agriculture, and 10% of North American agriculture is aimed at balancing the production of food with the production of other ecosystem services. Under this scenario, significant advances occur in the development of environmental technologies to increase production of services, create substitutes, and reduce harmful trade-offs.

Figure 13. MAIN DIRECT DRIVERS OF CHANGE IN BIODIVERSITY AND ECOSYSTEMS (CWG)

The cell color indicates impact of each driver on biodiversity in each type of ecosystem over the past 50–100 years. High impact means that over the last century the particular driver has significantly altered biodiversity in that biome; low impact indicates that it has had little influence on biodiversity in the biome. The arrows indicate the trend in the driver. Horizontal arrows indicate a continuation of the current level of impact; diagonal and vertical arrows indicate progressively increasing trends in impact. Thus, for example, if an ecosystem had experienced a very high impact of a particular driver in the past century (such as the impact of invasive species on islands), a horizontal arrow indicates that this very high impact is likely to continue. This Figure is based on expert opinion consistent with and based on the analysis of drivers of change in the various chapters of the assessment report of the MA Condition and Trends Working Group. The Figure presents global impacts and trends that may be different from those in specific regions.

		Habitat change	Climate change	Invasive species	Over-exploitation	Pollution (nitrogen, phosphorus)
Forest	Boreal	↗	↑	↗	→	↑
	Temperate	↘	↑	↑	→	↑
	Tropical	↑	↑	↑	↗	↑
Dryland	Temperate grassland	↗	↑	→	→	↑
	Mediterranean	↗	↑	↑	→	↑
	Tropical grassland and savanna	↗	↑	↑	→	↑
	Desert	→	↑	→	→	↑
Inland water		↑	↑	↑	→	↑
Coastal		↗	↑	↗	↗	↑
Marine		↑	↑	→	↗	↑
Island		→	↑	→	→	↑
Mountain		→	↑	→	→	↑
Polar		↗	↑	→	↗	↑

Driver's impact on biodiversity over the last century

- Low
- Moderate
- High
- Very high

Driver's current trends

- Decreasing impact
- Continuing impact
- Increasing impact
- Very rapid increase of the impact

Source: Millennium Ecosystem Assessment

increased risk of cancer and other chronic diseases from nitrates in drinking water, and increased risk of a variety of pulmonary and cardiac diseases from the production of fine particles in the atmosphere.

■ *Anthropogenic Climate Change*: Observed recent changes in climate, especially warmer regional temperatures, have already had significant impacts on biodiversity and ecosystems, including causing changes in species distributions, population sizes, the timing of reproduction or migration events, and an increase in the frequency of pest and disease outbreaks. Many coral reefs have undergone major, although often partially reversible, bleaching episodes when local sea surface temperatures have increased during one month by 0.5–1° Celsius above the average of the hottest months

By the end of the century, climate change and its impacts may be the dominant direct driver of biodiversity loss and changes in ecosystem services globally. The scenarios developed by the Intergovernmental Panel on Climate Change project an increase in global mean surface temperature of 2.0–6.4° Celsius above preindustrial levels by 2100, increased incidence of floods and droughts, and a rise in sea level of an additional 8–88 centimeters between 1990 and 2100. Harm to biodiversity will grow worldwide with increasing rates of change in climate and increasing absolute amounts of change. In contrast, some ecosystem services in some regions may initially be enhanced by projected changes in climate (such as increases in temperature or precipitation), and thus these regions may experience net benefits at low levels of climate change. As climate change becomes more severe, however, the harmful impacts on ecosystem services outweigh the benefits in most regions of the world. The balance of scientific evidence suggests that there will be a significant net harmful impact on ecosystem services worldwide if global mean surface temperature increases more than 2° Celsius above preindustrial levels or at rates greater than 0.2° Celsius per decade (*medium certainty*). There is a wide band of uncertainty in the amount of warming that would result from any stabilized greenhouse gas concentration, but based on IPCC projections this would require an eventual CO₂ stabilization level of less than 450 parts per million carbon dioxide (*medium certainty*).

Under all four MA scenarios, the projected changes in drivers result in significant growth in consumption of ecosystem services, continued loss of biodiversity, and further degradation of some ecosystem services. [5]

■ During the next 50 years, demand for food crops is projected to grow by 70–85% under the MA scenarios, and demand for water by between 30% and 85%. Water withdrawals in developing countries are projected to increase significantly under the scenarios, although these are projected to decline in industrial countries (*medium certainty*).


■ Food security is not achieved under the MA scenarios by 2050, and child malnutrition is not eradicated (and is projected to increase in some regions in some MA scenarios) despite increasing food supply and more diversified diets (*medium certainty*).

■ A deterioration of the services provided by freshwater resources (such as aquatic habitat, fish production, and water supply for households, industry, and agriculture) is found in the scenarios, particularly in those that are reactive to environmental problems (*medium certainty*).

■ Habitat loss and other ecosystem changes are projected to lead to a decline in local diversity of native species in all four MA scenarios by 2050 (*high certainty*). Globally, the equilibrium number of plant species is projected to be reduced by roughly 10–15% as the result of habitat loss alone over the period of 1970 to 2050 in the MA scenarios (*low certainty*), and other

Figure 14. GLOBAL TRENDS IN THE CREATION OF REACTIVE NITROGEN ON EARTH BY HUMAN ACTIVITY, WITH PROJECTION TO 2050 (R9 Fig 9.1)

Most of the reactive nitrogen produced by humans comes from manufacturing nitrogen for synthetic fertilizer and industrial use. Reactive nitrogen is also created as a by-product of fossil fuel combustion and by some (nitrogen-fixing) crops and trees in agroecosystems. The range of the natural rate of bacterial nitrogen fixation in natural terrestrial ecosystems (excluding fixation in agroecosystems) is shown for comparison. Human activity now produces approximately as much reactive nitrogen as natural processes do on the continents. (Note: The 2050 projection is included in the original study and is not based on MA Scenarios.)



factors such as overharvesting, invasive species, pollution, and climate change will further increase the rate of extinction.

The degradation of ecosystem services poses a significant barrier to the achievement of the Millennium Development Goals and the MDG targets for 2015. [3] The eight Millennium Development Goals adopted by the United Nations in 2000 aim to improve human well-being by reducing poverty, hunger, child and maternal mortality, by ensuring education for all, by controlling and managing diseases, by tackling gender disparity, by ensuring environmental sustainability, and by pursuing global partnerships. Under each of the MDGs, countries have agreed to targets to be achieved by 2015. Many of the regions facing the greatest challenges in achieving these targets coincide with regions facing the greatest problems of ecosystem degradation.

Although socioeconomic policy changes will play a primary role in achieving most of the MDGs, many of the targets (and goals) are unlikely to be achieved without significant improvement in management of ecosystems. The role of ecosystem changes in exacerbating poverty (Goal 1, Target 1) for some groups of people has been described already, and the goal of environmental sustainability, including access to safe drinking water (Goal 7, Targets 9, 10, and 11), cannot be achieved as long as most ecosystem services are being degraded. Progress toward three other MDGs is particularly dependent on sound ecosystem management:

■ **Hunger (Goal 1, Target 2):** All four MA scenarios project progress in the elimination of hunger but at rates far slower than needed to attain the internationally agreed target of halving, between 1990 and 2015, the share of people suffering from hunger. Moreover, the improvements are slowest in the regions in which the problems are greatest: South Asia and sub-Saharan Africa. Ecosystem condition, in particular climate, soil degradation, and water availability, influences progress toward this goal through its effect on crop yields as well as through impacts on the availability of wild sources of food.

■ **Child mortality (Goal 4):** Undernutrition is the underlying cause of a substantial proportion of all child deaths. Three of the MA scenarios project reductions in child undernourishment by 2050 of between 10% and 60% but undernourishment increases by 10% in *Order from Strength (low certainty)*. Child mortality is also strongly influenced by diseases associated with water quality. Diarrhea is one of the predominant causes of infant deaths worldwide. In sub-Saharan Africa, malaria additionally plays an important part in child mortality in many countries of the region.

■ **Disease (Goal 6):** In the more promising MA scenarios, progress toward Goal 6 is achieved, but under *Order from Strength* it is plausible that health and social conditions for the North and South could further diverge, exacerbating health problems in many low-income regions. Changes in ecosystems

influence the abundance of human pathogens such as malaria and cholera as well as the risk of emergence of new diseases. Malaria is responsible for 11% of the disease burden in Africa, and it is estimated that Africa's GDP could have been \$100 billion larger in 2000 (roughly a 25% increase) if malaria had been eliminated 35 years ago. The prevalence of the following infectious diseases is particularly strongly influenced by ecosystem change: malaria, schistosomiasis, lymphatic filariasis, Japanese encephalitis, dengue fever, leishmaniasis, Chagas disease, meningitis, cholera, West Nile virus, and Lyme disease.


Finding #4: *The challenge of reversing the degradation of ecosystems while meeting increasing demands for their services can be partially met under some scenarios that the MA considered, but these involve significant changes in policies, institutions, and practices that are not currently under way. Many options exist to conserve or enhance specific ecosystem services in ways that reduce negative trade-offs or that provide positive synergies with other ecosystem services.*

Three of the four MA scenarios show that significant changes in policies, institutions, and practices can mitigate many of the negative consequences of growing pressures on ecosystems, although the changes required are large and not currently under way. [5] All provisioning, regulating, and cultural ecosystem services are projected to be in worse condition in 2050 than they are today in only one of the four MA scenarios (*Order from Strength*). At least one of the three categories of services is in better condition in 2050 than in 2000 in the other three scenarios. (See Figure 15.) The scale of interventions that result in these positive outcomes are substantial and include significant investments in environmentally sound technology, active adaptive management, proactive action to address environmental problems before their full consequences are experienced, major investments in public goods (such as education and health), strong action to reduce socioeconomic disparities and eliminate poverty, and expanded capacity of people to manage ecosystems adaptively. However, even in scenarios where one or more categories of ecosystem services improve, biodiversity continues to be lost and thus the long-term sustainability of actions to mitigate degradation of ecosystem services is uncertain.

Past actions to slow or reverse the degradation of ecosystems have yielded significant benefits, but these improvements have generally not kept pace with growing pressures and demands. [8] Although most ecosystem services assessed in the MA are being degraded, the extent of that degradation would have been much greater without responses implemented in past decades. For example, more than 100,000 protected areas (including strictly protected areas such as national parks as well as areas managed for the sustainable use of natural ecosystems, including timber or wildlife harvest) covering about

Figure 15. NUMBER OF ECOSYSTEM SERVICES ENHANCED OR DEGRADED BY 2050 IN THE FOUR MA SCENARIOS

The Figure shows the net change in the number of ecosystem services enhanced or degraded in the MA scenarios in each category of services for industrial and developing countries expressed as a percentage of the total number of services evaluated in that category. Thus, 100% degradation means that all the services in the category were degraded in 2050 compared with 2000, while 50% improvement could mean that three out of six services were enhanced and the rest were unchanged or that four out of six were enhanced and one was degraded. The total number of services evaluated for each category was six provisioning services, nine regulating services, and five cultural services.



Source: Millennium Ecosystem Assessment

11.7% of the terrestrial surface have now been established, and these play an important role in the conservation of biodiversity and ecosystem services (although important gaps in the distribution of protected areas remain, particularly in marine and freshwater systems). Technological advances have also helped lessen the increase in pressure on ecosystems caused per unit increase in demand for ecosystem services.

Substitutes can be developed for some but not all ecosystem services, but the cost of substitutes is generally high, and substitutes may also have other negative environmental consequences. [8] For example, the substitution of vinyl, plastics, and metal for wood has contributed to relatively slow growth in global timber consumption in recent years. But while the availability of substitutes can reduce pressure on specific ecosystem services, they may not always have positive net benefits on the environment. Substitution of fuelwood by fossil fuels, for example, reduces pressure on forests and lowers indoor air pollution but it also increases net greenhouse gas emissions. Substitutes are also often costlier to provide than the original ecosystem services.

Ecosystem degradation can rarely be reversed without actions that address the negative effects or enhance the positive effects of one or more of the five indirect drivers of change: population change (including growth and migration), change in economic activity (including economic growth, disparities in wealth, and trade patterns), sociopolitical factors (including factors ranging from the presence of conflict to public participation in decision-making), cultural factors, and technological change. [4] Collectively these factors influence the level of production and consumption of ecosystem services and the sustainability of the production. Both economic growth and population growth lead to increased consumption of ecosystem services, although the harmful environmental impacts of any particular level of consumption depend on the efficiency of the technologies used to produce the service. Too often, actions to slow ecosystem degradation do not address these indirect drivers. For example, forest

management is influenced more strongly by actions outside the forest sector, such as trade policies and institutions, macroeconomic policies, and policies in other sectors such as agriculture, infrastructure, energy, and mining, than by those within it.

An effective set of responses to ensure the sustainable management of ecosystems must address the indirect and drivers just described and must overcome barriers related to [8]:

- Inappropriate institutional and governance arrangements, including the presence of corruption and weak systems of regulation and accountability.

- Market failures and the misalignment of economic incentives.

- Social and behavioral factors, including the lack of political and economic power of some groups (such as poor people, women, and indigenous peoples) that are particularly dependent on ecosystem services or harmed by their degradation.

- Underinvestment in the development and diffusion of technologies that could increase the efficiency of use of ecosystem services and could reduce the harmful impacts of various drivers of ecosystem change.

- Insufficient knowledge (as well as the poor use of existing knowledge) concerning ecosystem services and management, policy, technological, behavioral, and institutional responses that could enhance benefits from these services while conserving resources.

All these barriers are further compounded by weak human and institutional capacity related to the assessment and management of ecosystem services, underinvestment in the regulation and management of their use, lack of public awareness, and lack of awareness among decision-makers of both the threats posed by the degradation of ecosystem services and the opportunities that more sustainable management of ecosystems could provide.

The MA assessed 74 response options for ecosystem services, integrated ecosystem management, conservation and sustainable use of biodiversity, and climate change. Many of these options hold significant promise for overcoming these barriers and conserving or sustainably enhancing the supply of ecosystem services. Promising options for specific sectors are shown in Box 2, while cross-cutting responses addressing key obstacles are described in the remainder of this section.

Institutions and Governance

Changes in institutional and environmental governance frameworks are sometimes required to create the enabling conditions for effective management of ecosystems, while in other cases existing institutions could meet these needs but face significant barriers. [8] Many existing institutions at both the global and the national level have the mandate to address the degradation of ecosystem services but face a variety of challenges in doing so related in part to the need for greater cooperation across sectors and the need for coordinated responses at multiple scales.

However, since a number of the issues identified in this assessment are recent concerns and were not specifically taken into account in the design of today's institutions, changes in existing institutions and the development of new ones may sometimes be needed, particularly at the national scale.

In particular, existing national and global institutions are not well designed to deal with the management of common pool resources, a characteristic of many ecosystem services. Issues of ownership and access to resources, rights to participation in decision-making, and regulation of particular types of resource use or discharge of wastes can strongly influence the sustainability of ecosystem management and are fundamental determinants of who wins and loses from changes in ecosystems. Corruption, a major obstacle to effective management of ecosystems, also stems from weak systems of regulation and accountability.

Promising interventions include:

- *Integration of ecosystem management goals within other sectors and within broader development planning frameworks.* The most important public policy decisions affecting ecosystems are often made by agencies and in policy arenas other than those charged with protecting ecosystems. For example, the Poverty Reduction Strategies prepared by developing-country governments for the World Bank and other institutions strongly shape national development priorities, but in general these have not taken into account the importance of ecosystems to improving the basic human capabilities of the poorest.

- *Increased coordination among multilateral environmental agreements and between environmental agreements and other international economic and social institutions.* International agreements are indispensable for addressing ecosystem-related concerns that span national boundaries, but numerous obstacles weaken their current effectiveness. Steps are now being taken to increase the coordination among these mechanisms, and this could help to broaden the focus of the array of instruments. However, coordination is also needed between the multilateral environmental agreements and more politically powerful international institutions, such as economic and trade agreements, to ensure that they are not acting at cross-purposes. And implementation of these agreements needs to be coordinated among relevant institutions and sectors at the national level.

- *Increased transparency and accountability of government and private-sector performance on decisions that have an impact on ecosystems, including through greater involvement of concerned stakeholders in decision-making.* Laws, policies, institutions, and markets that have been shaped through public participation in decision-making are more likely to be effective and perceived as just. Stakeholder participation also contributes to the decision-making process because it allows a better understanding of impacts and vulnerability, the distribution of costs and benefits associated with trade-offs, and the identification of a broader range of response options that are available in a specific context. And stakeholder involvement and transparency of decision-making can increase accountability and reduce corruption.

Box 2. EXAMPLES OF PROMISING AND EFFECTIVE RESPONSES FOR SPECIFIC SECTORS

Illustrative examples of response options specific to particular sectors judged to be promising or effective are listed below. (See Appendix B.) A response is considered effective when it enhances the target ecosystem services and contributes to human well-being without significant harm to other services or harmful impacts on other groups of people. A response is considered promising if it does not have a long track record to assess but appears likely to succeed or if there are known ways of modifying the response so that it can become effective.

Agriculture

- Removal of production subsidies that have adverse economic, social, and environmental effects.
- Investment in, and diffusion of, agricultural science and technology that can sustain the necessary increase of food supply without harmful tradeoffs involving excessive use of water, nutrients, or pesticides.
- Use of response policies that recognize the role of women in the production and use of food and that are designed to empower

women and ensure access to and control of resources necessary for food security.

- Application of a mix of regulatory and incentive- and market-based mechanisms to reduce overuse of nutrients.

Fisheries and Aquaculture

- Reduction of marine fishing capacity.
- Strict regulation of marine fisheries both regarding the establishment and implementation of quotas and steps to address unreported and unregulated harvest. Individual transferable quotas may be appropriate in some cases, particularly for cold water, single species fisheries.
- Establishment of appropriate regulatory systems to reduce the detrimental environmental impacts of aquaculture.
- Establishment of marine protected areas including flexible no-take zones.

Water

- Payments for ecosystem services provided by watersheds.
- Improved allocation of rights to freshwater resources to align incentives with conservation needs.

- Increased transparency of information regarding water management and improved representation of marginalized stakeholders.
- Development of water markets.
- Increased emphasis on the use of the natural environment and measures other than dams and levees for flood control.
- Investment in science and technology to increase the efficiency of water use in agriculture.

Forestry

- Integration of agreed sustainable forest management practices in financial institutions, trade rules, global environment programs, and global security decision-making.
- Empowerment of local communities in support of initiatives for sustainable use of forest products; these initiatives are collectively more significant than efforts led by governments or international processes but require their support to spread.
- Reform of forest governance and development of country-led, strategically focused national forest programs negotiated by stakeholders.

Economics and Incentives

Economic and financial interventions provide powerful instruments to regulate the use of ecosystem goods and services. [8] Because many ecosystem services are not traded in markets, markets fail to provide appropriate signals that might otherwise contribute to the efficient allocation and sustainable use of the services. A wide range of opportunities exists to influence human behavior to address this challenge in the form of economic and financial instruments. However, market mechanisms and most economic instruments can only work effectively if supporting institutions are in place, and thus there is a need to build institutional capacity to enable more widespread use of these mechanisms.

Promising interventions include:

- *Elimination of subsidies that promote excessive use of ecosystem services (and, where possible, transfer of these subsidies to payments for non-marketed ecosystem services).* Government subsidies paid to the agricultural sectors of OECD countries between 2001 and 2003 averaged over \$324 billion annually, or one third the global value of agricultural products in 2000. A significant proportion of this total involved production subsidies that led to greater

food production in industrial countries than the global market conditions warranted, promoted overuse of fertilizers and pesticides in those countries, and reduced the profitability of agriculture in developing countries. Many countries outside the OECD also have inappropriate input and production subsidies, and inappropriate subsidies are common in other sectors such as water, fisheries, and forestry. Although removal of perverse subsidies will produce net benefits, it will not be without costs. Compensatory mechanisms may be needed for poor people who are adversely affected by the removal of subsidies, and removal of agricultural subsidies within the OECD would need to be accompanied by actions designed to minimize adverse impacts on ecosystem services in developing countries.

- *Greater use of economic instruments and market-based approaches in the management of ecosystem services.* These include:
 - Taxes or user fees for activities with “external” costs (tradeoffs not accounted for in the market). Examples include taxes on excessive application of nutrients or ecotourism user fees.

- Creation of markets, including through cap-and-trade systems. One of the most rapidly growing markets related to ecosystem services is the carbon market. Approximately 64 million tons of carbon dioxide equivalent were exchanged through projects from January to May 2004, nearly as much as during all of 2003. The value of carbon trades in 2003 was approximately \$300 million. About one quarter of the trades involved investment in ecosystem services (hydropower or biomass). It is *speculated* that this market may grow to \$10 billion to \$44 billion by 2010. The creation of a market in the form of a nutrient trading system may also be a low-cost way to reduce excessive nutrient loading in the United States.
- Payment for ecosystem services. For example, in 1996 Costa Rica established a nationwide system of conservation payments to induce landowners to provide ecosystem services. Under this program, Costa Rica brokers contracts between international and domestic “buyers” and local “sellers” of sequestered carbon, biodiversity, watershed services, and scenic beauty. Another innovative conservation financing mechanism is “biodiversity offsets,” whereby developers pay for conservation activities as compensation for unavoidable harm that a project causes to biodiversity.
- Mechanisms to enable consumer preferences to be expressed through markets. For example, current certification schemes for sustainable fisheries and forest practices provide people with the opportunity to promote sustainability through their consumer choices.

Social and Behavioral Responses

Social and behavioral responses—including population policy, public education, civil society actions, and empowerment of communities, women, and youth—can be instrumental in responding to the problem of ecosystem degradation. [8] These are generally interventions that stakeholders initiate and execute through exercising their procedural or democratic rights in efforts to improve ecosystems and human well-being.

Promising interventions include:

- *Measures to reduce aggregate consumption of unsustainably managed ecosystem services.* The choices about what individuals consume and how much are influenced not just by considerations of price but also by behavioral factors related to culture, ethics, and values. Behavioral changes that could reduce demand for degraded ecosystem services can be encouraged through actions by governments (such as education and public awareness programs or the promotion of demand-side management), industry (commitments to use raw materials that are from sources certified as being sustainable, for example, or improved product labeling), and civil society (through raising public awareness). Efforts to reduce aggregate consumption, however, must sometimes incorporate measures to increase the access to and consumption of those same ecosystem services by specific groups such as poor people.

- *Communication and education.* Improved communication and education are essential to achieve the objectives of environmental conventions and the Johannesburg Plan of Implementation as well as the sustainable management of natural resources more generally. Both the public and decision-makers can benefit from education concerning ecosystems and human well-being, but education more generally provides tremendous social benefits that can help address many drivers of ecosystem degradation. While the importance of communication and education is well recognized, providing the human and financial resources to undertake effective work is a continuing problem.

- *Empowerment of groups particularly dependent on ecosystem services or affected by their degradation, including women, indigenous peoples, and young people.* Despite women’s knowledge about the environment and the potential they possess, their participation in decision-making has often been restricted by economic, social, and cultural structures. Young people are also key stakeholders in that they will experience the longer-term consequences of decisions made today concerning ecosystem services. Indigenous control of traditional homelands can sometimes have environmental benefits, although the primary justification continues to be based on human and cultural rights.

Technological Responses

Given the growing demands for ecosystem services and other increased pressures on ecosystems, the development and diffusion of technologies designed to increase the efficiency of resource use or reduce the impacts of drivers such as climate change and nutrient loading are essential. [8] Technological change has been essential for meeting growing demands for some ecosystem services, and technology holds considerable promise to help meet future growth in demand. Technologies already exist for reduction of nutrient pollution at reasonable costs—including technologies to reduce point source emissions, changes in crop management practices, and precision farming techniques to help control the application of fertilizers to a field, for example—but new policies are needed for these tools to be applied on a sufficient scale to slow and ultimately reverse the increase in nutrient loading (even while increasing nutrient application in regions such as sub-Saharan Africa where too little fertilizer is being applied). However, negative impacts on ecosystems and human well-being have sometimes resulted from new technologies, and thus careful assessment is needed prior to their introduction.

Promising interventions include:

- *Promotion of technologies that enable increased crop yields without harmful impacts related to water, nutrient, and pesticide use.* Agricultural expansion will continue to be one of the major drivers of biodiversity loss well into the twenty-first century. Development, assessment, and diffusion of technologies that could increase the production of food per unit area sustainably without harmful trade-offs related to excessive consumption of water or use of nutrients or pesticides would significantly lessen pressure on other ecosystem services.



■ *Restoration of ecosystem services.* Ecosystem restoration activities are now common in many countries. Ecosystems with some features of the ones that were present before conversion can often be established and can provide some of the original ecosystem services. However, the cost of restoration is generally extremely high compared with the cost of preventing the degradation of the ecosystem. Not all services can be restored, and heavily degraded services may require considerable time for restoration.

■ *Promotion of technologies to increase energy efficiency and reduce greenhouse gas emissions.* Significant reductions in net greenhouse gas emissions are technically feasible due to an extensive array of technologies in the energy supply, energy demand, and waste management sectors. Reducing projected emissions will require a portfolio of energy production technologies ranging from fuel switching (coal/oil to gas) and increased power plant efficiency to increased use of renewable energy technologies, complemented by more efficient use of energy in the transportation, buildings, and industry sectors. It will also involve the development and implementation of supporting institutions and policies to overcome barriers to the diffusion of these technologies into the marketplace, increased public and private-sector funding for research and development, and effective technology transfer.

Knowledge Responses

Effective management of ecosystems is constrained both by the lack of knowledge and information about different aspects of ecosystems and by the failure to use adequately the information that does exist in support of management decisions.

[8, 9] In most regions, for example, relatively limited information exists about the status and economic value of most ecosystem

services, and their depletion is rarely tracked in national economic accounts. Basic global data on the extent and trend in different types of ecosystems and land use are surprisingly scarce. Models used to project future environmental and economic conditions have limited capability of incorporating ecological “feedbacks,” including nonlinear changes in ecosystems, as well as behavioral feedbacks such as learning that may take place through adaptive management of ecosystems.

At the same time, decision-makers do not use all of the relevant information that is available. This is due in part to institutional failures that prevent existing policy-relevant scientific information from being made available to decision-makers and in part to the failure to incorporate other forms of knowledge and information (such as traditional knowledge and practitioners’ knowledge) that are often of considerable value for ecosystem management.

Promising interventions include:

■ *Incorporation of nonmarket values of ecosystems in resource management and investment decisions.* Most resource management and investment decisions are strongly influenced by considerations of the monetary costs and benefits of alternative policy choices. Decisions can be improved if they are informed by the total economic value of alternative management options and involve deliberative mechanisms that bring to bear noneconomic considerations as well.

■ *Use of all relevant forms of knowledge and information in assessments and decision-making, including traditional and practitioners' knowledge.* Effective management of ecosystems typically requires “place-based” knowledge—that is, information about the specific characteristics and history of an ecosystem. Traditional knowledge or practitioners' knowledge held by local resource managers can often be of considerable value in resource management, but it is too rarely incorporated into decision-making processes and indeed is often inappropriately dismissed.

■ *Enhancing and sustaining human and institutional capacity for assessing the consequences of ecosystem change for human well-being and acting on such assessments.* Greater technical capacity is



needed for agriculture, forest, and fisheries management. But the capacity that exists for these sectors, as limited as it is in many countries, is still vastly greater than the capacity for effective management of other ecosystem services.

A variety of frameworks and methods can be used to make better decisions in the face of uncertainties in data, prediction, context, and scale. Active adaptive management can be a particularly valuable tool for reducing uncertainty about ecosystem management decisions. [8] Commonly used decision-support methods include cost-benefit analysis, risk assessment, multicriteria analysis, the precautionary principle, and vulnerability analysis. Scenarios also provide one means to cope with many aspects of uncertainty, but our limited understanding of ecological systems and human responses shrouds any individual scenario in its own characteristic uncertainty. Active adaptive management is a tool that can be particularly valuable given the high levels of uncertainty surrounding coupled socioecological systems. This involves the design of management programs to test hypotheses about how components of an ecosystem function and interact, thereby reducing uncertainty about the system more rapidly than would otherwise occur.

Sufficient information exists concerning the drivers of change in ecosystems, the consequences of changes in ecosystem services for human well-being, and the merits of various response options to enhance decision-making in support of sustainable development at all scales. However, many research needs and information gaps were identified in this assessment, and actions to address those needs could yield substantial benefits in the form of improved information for policy and action. [9] Due to gaps in data and knowledge, this assessment was unable to answer fully a number of questions posed by its users. Some of these gaps resulted from weaknesses in monitoring systems related to ecosystem services and their linkages with human well-being. In other cases, the assessment revealed significant needs for further research, such as the need to improve understanding of nonlinear changes in ecosystems and of the economic value of alternative management options. Investments in improved monitoring and research, combined with additional assessments of ecosystem services in different nations and regions, would significantly enhance the utility of any future global assessment of the consequences of ecosystem change for human well-being.